

December 14, 2012

Mr. David Ferris  
Division of Wastewater Management  
Department of Environmental Protection  
1 Winter Street  
Boston, MA 02108

Mr. Stephen Perkins  
U.S. Environmental Protection Agency  
Water Enforcement  
OES4-SMR  
5 Post Office Square, Suite 100  
Boston, MA 02109-3912

RE: Massachusetts Water Resources Authority  
Permit Number MA 0103284  
MWRA Contingency Plan Threshold Exceedance: Infaunal diversity for August 2012

Dear Mr. Ferris and Mr. Perkins:

In its outfall ambient monitoring program, MWRA measures the numbers and types of infaunal organisms living in sediments in western Massachusetts Bay in the vicinity of the outfall and in reference areas. In 2012, samples from 11 nearfield stations were collected in August. Four separate measures of infaunal community diversity in nearfield sediments are in the Contingency Plan. Other measures of the health of sediment communities in the Contingency Plan are the depth of oxygen penetration into nearfield sediments and the abundance of six opportunistic species.<sup>1</sup>

On December 10, 2012, MWRA completed its data quality reviews and calculations of Contingency Plan threshold values for nearfield infaunal diversity and for the abundance of opportunists for its August 2012 monitoring. Those results and results for depth of oxygen penetration (apparent RPD) are in Table 1. Two of the diversity parameters calculated, Shannon-Wiener  $H'$  and Pielou's  $J'$ , were slightly higher (more diverse) than their upper thresholds. Those results are a "Caution Level" threshold exceedance requiring regulatory and public notification. This letter constitutes that notification. MWRA is evaluating the 2012 data in more detail, however, there are currently no indications that this threshold exceedance was influenced by the outfall discharge, nor are there any indications of a decline in the health of sediment communities.

---

<sup>1</sup> *Massachusetts Water Resources Authority Contingency Plan Revision 1*. 2001. Report ms-071.  
<http://www.mwra.state.ma.us/harbor/enquad/trlist.html>

The two other diversity parameters calculated for August 2012 samples, log-series alpha and total species richness, were within their threshold ranges. The six opportunistic taxa made up less than 1% of the animals collected from nearfield sediments (the Caution Level is 10%). The apparent Redox Potential Discontinuity (RPD) was 3.44 cm, more than twice as deep as the minimum threshold.

**Table 1. Infaunal monitoring threshold results, August 2012 Samples**

Parameter	Threshold range		Result	Exceedance?
	Low	High		
Total species	43.0	81.9	73.45	No
Log-series Alpha	9.42	15.8	15.68	No
Shannon-Weiner H'	3.37	3.99	4.36	Yes, Caution Level
Pielou's J'	0.57	0.67	0.70	Yes, Caution Level
Apparent RPD	1.18	NA	3.44	No
Percent opportunists	10% (Caution) 25% (Warning)		0.16%	No

Similar exceedances of the same two diversity indices were observed in 2010 and 2011 ([http://www.mwra.state.ma.us/harbor/pdf/20110107amx\\_diversity.pdf](http://www.mwra.state.ma.us/harbor/pdf/20110107amx_diversity.pdf) and [http://www.mwra.state.ma.us/harbor/pdf/20111215amx\\_diversity.pdf](http://www.mwra.state.ma.us/harbor/pdf/20111215amx_diversity.pdf)). The 2010 exceedance was evaluated and discussed at the June 2011 meeting of EPA's Outfall Monitoring Science Advisory Panel (OMSAP), with details of the evaluation included in the 2010 Outfall Benthic Monitoring Report.<sup>2</sup> The 2011 exceedance was evaluated in the 2011 Outfall Benthic Monitoring Report<sup>3</sup>. Those evaluations found that the 2010 and 2011 exceedances probably represented natural fluctuations in the infaunal communities, and were not influenced by the outfall.

Thus far, the 2012 data support the same conclusion.

Given the strong year-to-year similarity normally observed in infaunal communities during MWRA's monitoring, it is not surprising that similar exceedances in the same parameters have been observed in each of the past 3 years.

## Background

Sediment monitoring in the vicinity of the outfall was designed to address questions about potential effects of the relocated discharge on the offshore seafloor including eutrophication and related low levels of dissolved oxygen, accumulation of toxic contaminants in depositional areas, and smothering of animals by particulate matter.

*Monitoring design.* Soft sediment samples are collected for the identification and enumeration of the benthic infaunal community. Samples are also collected for the analysis of grain size, total organic carbon (TOC), and the effluent solids tracer bacterium *Clostridium perfringens* spores. Sediment profile imaging, done with a camera, provides a rapid visual assessment of sediment quality.

<sup>2</sup> Maciolek NJ, Dahlen DT, Diaz RJ, Hecker B. 2011. Outfall benthic monitoring report: 2010 results. Boston: Massachusetts Water Resources Authority. Report 2011-14. <http://www.mwra.state.ma.us/harbor/enquad/trlist.html>

<sup>3</sup> Nestler EC, Diaz RJ, Hecker B, Pembroke AE. 2012. Outfall benthic monitoring report: 2011 results. Boston: Massachusetts Water Resources Authority. Report 2012-08. 38 pages plus appendices. <http://www.mwra.state.ma.us/harbor/enquad/pdf/2012-08.pdf>

The August 2012 sediment sampling was conducted according to MWRA's revised ambient monitoring plan<sup>4</sup> and included sediment profile imaging at 23 nearfield stations (Figure 1) and infaunal sampling and sampling for sediment grain size, total organic carbon, and *Clostridium perfringens* spores at the 11 nearfield and 2 farfield stations shown in Figure 2.

*Infaunal thresholds.* Infaunal thresholds are triggered by nearfield results outside of those found during baseline monitoring in 1992-2000. Low levels of enrichment can fertilize communities, increasing diversity, while higher levels cause stress, leading to lower species richness, lower evenness, and lower biodiversity and increases in the abundance of opportunistic species, which are able to persist in degraded environments.<sup>5</sup> The Contingency Plan defines the Caution Level as "appreciable change" in the following six infaunal thresholds:

Four biodiversity thresholds. *Total species* is the number of species identified and is the simplest measure of how many species are present. *Log-series alpha* is another measure of species richness with some theoretical advantages over total species. *Pielou's J'* measures how evenly individuals are distributed among species.<sup>6</sup> *Shannon-Wiener H'* is a commonly used diversity index affected by both species richness and evenness. The infaunal thresholds were chosen to be triggered if the nearfield mean diversity for a discharge year was *lower* than the 2.5<sup>th</sup> percentile of the distribution of baseline means, or *higher* than the 97.5<sup>th</sup> percentile<sup>7</sup>.

The August 2012 results for Shannon-Wiener H' and Pielou's J' are slightly higher than the upper threshold (Table 1) triggering the exceedance.

Percent opportunistic species. Increased deposition of organic solids can increase abundances of opportunistic species. The Caution Level threshold for the abundance of six opportunistic species<sup>8</sup> was set at 10% of total nearfield infaunal abundances, roughly 20% of their abundance in Boston Harbor during 1991-1998.

Depth of penetration of oxygen. The depth below the sediment-water interface where sediments change from light-colored, oxygenated mud to black anoxic mud is called the depth of the apparent redox potential discontinuity (RPD). Microbial breakdown of organic material uses up oxygen, decreasing the thickness of the RPD, so deeper, bigger RPDs are considered healthier. The threshold was set at half the average RPD measured during baseline.

---

<sup>4</sup> MWRA. 2010. Massachusetts Water Resources Authority effluent outfall ambient monitoring plan Revision 2. July 2010. Boston: Massachusetts Water Resources Authority. Report 2010-04. 107 p.

<sup>5</sup> In addition to the thresholds discussed here, MWRA tests the average concentrations of 26 organic and heavy metal contaminants in nearfield sediments. Samples for contaminants were collected in August 2011 (all results were below the threshold concentrations), and samples will next be collected in August 2014.

<sup>6</sup> Communities with a similar number of individuals from a number of different species, (for example a temperate forest with similar numbers of oak, maple, elm, etc.) are more diverse than ecosystems with the same number of species but with the overwhelming majority made up of one species (for example a wheat field where 99% of the plants are wheat but a few different types of weeds are growing among the crop).

<sup>7</sup> Infaunal thresholds were recalculated using the baseline data collected at the 11 stations now sampled under the revised ambient monitoring design.

<sup>8</sup> The six species include the polychaete worms *Capitella capitata* species complex, *Streblospio benedicti*, and *Polydora cornuta*, the amphipod crustaceans *Ampelisca abdita* and *A. vadorum*, and the bivalve mollusk *Mulinia lateralis*.




Figure 1. Map of sediment profile imaging stations.





Figure 2. Map of infaunal community monitoring stations sampled in 2011 and 2012.

## 2012 results


It is important to note that the evaluation of the 2012 infaunal results is just beginning, and will continue in the weeks and months to come.

*Sediment conditions.* All indications available at this time are that conditions in nearfield sediments in 2012 were normal. Preliminary review of the sediment profile image results from all 23 nearfield stations in August 2012 suggested that conditions were similar to observations made in 2011, with deeper oxygen penetration into the sediments than seen since 2007 (Figure 3) and evidence for robust biological reworking of sediments at many sites (for example burrows and subsurface feeding voids).

Total organic carbon and grain size measured at the nearfield stations in 2012 were also well within baseline and outfall discharge ranges for these stations, as was the effluent solids tracer, spores of the bacterium *Clostridium perfringens* (Figure 4).



**Figure 3 . Depth of the average redox potential discontinuity (RPD) in nearfield sediments as measured from sediment profile images, 1992-2012. RPDs in 2012 are among the deepest measured during discharge monitoring, indicating healthy sediment oxygenation.**



**Figure 4. Scatterplot of *Clostridium perfringens* spores and percent silt/clay in sediments, 1992-2012. 2012 data are well within previous measurements. Gray dots are baseline period data, green triangles are outfall discharge period data, and blue diamonds are 2012 data.**

So far, MWRA’s review of the 2012 nearfield infaunal results provides no indications that the slightly elevated Pielou’s  $J'$  and Shannon-Wiener  $H'$  diversities (Figure 5) reflect a response to the outfall discharge. Since both of the diversity indices sensitive to species richness were high but not extremely high in 2011 (Figure 6) it is likely that increased species evenness is responsible for the threshold exceedance, both for  $J'$ , a measure of evenness, and  $H'$ , affected both by species richness and by evenness in samples. This was also observed in MWRA’s evaluations of the 2010 and 2011 threshold exceedances. Infaunal densities in 2012 were somewhat higher than observed in 2010 and 2011, averaging about 44,000 animals/m<sup>2</sup>. High evenness in abundant, species-rich samples is not typical of stressed communities, therefore it is likely that the results observed represent a natural fluctuation in the nearfield infaunal communities. The observation of continued very low abundances of opportunistic taxa (Figure 7) also supports this.

Evaluation of the 2012 sediment nearfield monitoring data is continuing.




Figure 5. Pielou's  $J'$  evenness (top) and Shannon-Wiener  $H'$  diversity (bottom) measured at current nearfield monitoring stations, 1992-2003 and 2011-2012. Data from 2004 through 2010 are the averages for the odd and even-year stations sampled then. Blue points are baseline period data, and green triangles reflect outfall discharge period data. The threshold levels varied slightly through the monitoring period, for simplicity only the current thresholds are shown (dotted lines are the current upper and lower thresholds).  $H'$  diversity in 2004 ( $H' = 4.08$ ) did not exceed the threshold (4.14) for even-year stations applicable in 2004.






Figure 6. Log-series alpha and total species richness measured at nearfield monitoring stations, 1992-2012. Blue points are baseline period data, and green triangles reflect outfall discharge period data. The threshold levels varied slightly through the monitoring period, for simplicity only the current thresholds are shown (dotted lines are the current upper and lower thresholds).




Figure 7. Abundance of opportunist taxa measured nearfield monitoring stations, 1992-2012. The dotted line reflects the 10% caution threshold. The 25% warning threshold is not plotted. Blue points are baseline period data, and green triangles reflect outfall discharge period data. The thresholds have remained the same throughout the monitoring.

MWRA will discuss this threshold exceedance at the next Outfall Monitoring Science Advisory Panel meeting.

If you have questions or need additional information, please feel free to call Dr. Andrea Rex at (617) 788-4940.

Sincerely,

Michael J. Hornbrook  
 Chief Operating Officer

Cc:

**Environmental Protection Agency, Region I**

Matthew Liebman  
Todd Borci

**National Marine Fisheries Service**

Daniel Morris

**Stellwagen Bank National Marine Sanctuary**

Craig MacDonald

**MA EOEEA**

Kathy Baskin

**MA Department of Environmental Protection**

Cathy Vakalopoulos  
David Ferris

**MA Division of Marine Fisheries**

Jack Schwartz

**Cape Cod Commission**

Tom Cambareri

**Outfall Monitoring Science Advisory Panel**

Andrew Solow  
Robert Beardsley  
Norb Jaworski  
Robert Kenney  
Judy Pederson  
Michael Shiaris  
James Shine  
Juanita Urban-Rich

**Public Interest Advisory Committee**

Patty Foley

**Hyannis Library**

Ann-Louise Harries

**MWRA Library**

Elizabeth Steele